

A Simple Approach to Staggered Difference-in-Differences in the Presence of Spillovers

Mario Fiorini * Wooyong Lee † Gregor Pfeifer ‡

March 14, 2024

Abstract

We establish identifying assumptions and estimation procedures for the ATT in a Difference-in-Differences setting with staggered treatment adoption in the presence of spillovers. We show that the ATT can be estimated by a simple TWFE method that extends the approach of [Wooldridge \[2022\]](#)'s fully interacted regression model. We broaden our framework to the non-linear case of count data, offering estimation of the ATT by a simple TWFE Poisson model, and we revisit a corresponding application from the crime literature. Monte Carlo simulations show that our estimator performs competitively.

Keywords: Difference-in-Differences, staggered treatment adoption, spillovers, (non-)linear models.

*(corresponding author): University of Technology Sydney (mario.fiorini@uts.edu.au).

†University of Technology Sydney (wooyong.lee.econ@gmail.com)

‡University of Sydney, CESifo, and IZA (gregor.pfeifer@sydney.edu.au).

1 Introduction

The Difference-in-Differences (DiD) literature, particularly the one concerned with staggered treatment adoption, has experienced significant advances in the last few years, and papers by [Roth et al. \[2023\]](#) and [de Chaisemartin and D’Haultfoeuille \[2021\]](#) have summarized these developments. Within this array of advances, one area still understudied is the one linked to spillovers—implying that the Stable Unit Treatment Value Assumption (SUTVA) assumption does not hold. However, as [Roth et al. \[2023\]](#) point out, spillover effects may be important in many economic applications, such as when a policy in one area affects neighboring areas, or when individuals are connected in a network.¹ Our work contributes to this area and links two active DiD literature strands.

The first one focuses on estimation issues under staggered adoption and heterogeneous treatment effects across units and time. [Borusyak et al. \[2021\]](#), [de Chaisemartin and D’Haultfoeuille \[2020\]](#), [Callaway and Sant’Anna \[2020\]](#), [Goodman-Bacon \[2021\]](#), [Sun and Abraham \[2020\]](#) and [Wooldridge \[2022\]](#) highlight that the two-way fixed effect (TWFE) regression estimator may be biased for the average treatment effect on the treated (ATT), to the extreme of showing the opposite sign. The authors suggest alternative estimators that account for the variation in treatment timing, thereby providing a consistent estimator for the ATT. We contribute to this literature by extending it to the case of spillovers in both linear and non-linear models.

The second strand studies the identification of average treatment effects in the presence of spillovers. [Berg et al. \[2021\]](#), [Butts \[2023\]](#), [Clarke \[2017\]](#), and [Huber and Steinmayr \[2021\]](#) highlight two main challenges for identification of the ATT if the treatment also impacts units that are not formally treated. First, untreated units are no longer valid controls. So far, proposed solutions mostly centre around ruling out spillovers for a given group of units, often based on some spatial distance, allowing the researcher to use this latter group as a control. Alternatively, if sufficient information exists, one can parametrize how units are exposed to spillovers. Second, multiple definitions of the ATT are possible in the presence of spillovers. This is because a unit’s treatment can lead to changes to its own outcome, but also to other units’ outcomes. In this case, the researcher might be interested in examining the former effect, summarized by the ATT without interference (i.e., the one identified under SUTVA), or in a broader definition of the ATT that also accounts for the latter effect. Here, we contribute to this literature by providing conditions that allow for the identification of the ATT without interference, despite the presence of spillovers. Our setting also departs from this literature since we focus on the more complex staggered treatment adoption, which has the potential for

¹As another example, [Minton and Mulligan \[2024\]](#) use price theory to demonstrate that when treated and control units are in the same market, control units are indirectly affected by the treatment.

cumulative spillovers. Nevertheless, our results also apply to the simpler simultaneous treatment case.

Specifically regarding contributions, we first establish the identifying assumptions for the ATT without interference given a staggered DiD setting in the presence of spillovers. We show that aside from the canonical i) treatment irreversibility, ii) no-anticipation, and iii) parallel trends assumptions, identification requires that once a unit receives treatment, it is no longer influenced by spillover effects. This means the unit forfeits any spillovers it may have previously received and remains unaffected by spillovers from subsequently treated groups. This assumption also unifies the multiple definitions of the ATT, because they are the same with or without spillovers, simplifying policy evaluation and joining with the definition of ATT under SUTVA. We also assume that a set of never-treated units is not exposed to spillovers, in line with the existing literature. The combination of these assumptions allows for the identification of the ATT. Below, we argue that such a scenario applies to many contexts. Differently from [Butts \[2023\]](#), who is closest to our work, we directly focus on the staggered treatment scenario and, importantly, provide assumptions for the identification of the ATT without interference.²

Our second contribution regards estimation. We show that the extended TWFE model approach of [Wooldridge \[2022\]](#), which is numerically equivalent to the imputation-based approach of [Borusyak et al. \[2021\]](#), can be used to account for spillovers. Furthermore, we discuss identification and estimation in the non-linear case of count data, broadening the range of applications where our approach can be applied to. For our empirical application, we revisit [Gonzalez-Navarro \[2013\]](#), who studied the effects of installing a stolen vehicle recovery device on car theft. Since car theft is a count variable, we implement the non-linear Poisson DiD adjusted for spillovers. Our correction leads to a larger effect of the policy relative to the original contribution’s specification.

Finally, we perform a Monte Carlo analysis, highlighting the bias-variance trade-off implicit in the correction for staggered treatment and spillovers. Identification of time and group fixed effects can neither rely on the already treated units due to heterogeneous treatment effects, nor on the untreated units potentially exposed to spillovers. However, the benefit of excluding such units from estimation can be small if treatment effects are relatively homogeneous and if spillovers are small, while costing the researcher precision. We compare the traditional TWFE estimator, which ignores both staggered adoption and spillovers, the [Wooldridge \[2022\]](#) estimator, which accounts for staggered adoption but not for spillovers, and our estimator, which corrects for both. We do so under different sample sizes, degrees of staggered treatment, and degrees of spillovers, showing that our estimator performs competitively in many settings.

²[Butts \[2023\]](#) is concerned with establishing identification of the sum of direct and spillover effects.

The remainder of the paper is organized as follows. Section 2 provides intuition alongside two motivating examples, after which Section 3 lays out the formal DiD setup with staggered treatment adoption. Section 4 establishes conditions for identifying the ATT, while Section 5 discusses estimation and inference considering the formerly established assumptions. Section 6 extends our model to the non-linear case, and Section 7 discusses a corresponding application. Section 8 provides Monte Carlo simulations, and Section 9 concludes.

2 Intuition and Motivating Examples

To illustrate our setting, consider panel data of units divided into three groups, A, B, and Z, observed over three periods, 1, 2, and 3. The groups are distinguished by the timing of their treatment. Group A is treated in periods 2 and 3, group B in period 3, and group Z remains untreated throughout. Each group consists of two units, denoted by a, a', b, b', z, z' . We use the indices i and j to refer to any unit within these groups. Figure 1 illustrates the potential treatment and spillover mechanisms in this setting, focusing on periods 2 and 3. Solid lines indicate treatment effects under no interference (β_{it}), equivalent to the conventional definition of the treatment effect under SUTVA. We call them direct effects. Dotted lines indicate spillovers from a treated unit j to a treated unit i (γ_{it}^j), and dashed ones represent spillovers from a treated unit j to an untreated unit i (η_{it}^j). The figure highlights a key challenge introduced by the presence of spillovers: there are no valid control units. This is further complicated by the presence of multiple direct effects, which is a feature of staggered adoption design. Under SUTVA, only the direct effects represented by solid lines would exist.

We can visualize possible data patterns in this setting using a simple parametric model. Suppose that the outcome of interest is deterministic and given by:

$$Y_{it} = 1 + \delta_t + \beta_{it} \cdot D_{it} + D_{it} \cdot \sum_{j \neq i} \gamma_{it}^j \cdot D_{jt} + (1 - D_{it}) \cdot \sum_{j \neq i} \eta_{it}^j \cdot D_{jt}, \quad (1)$$

where D_{it} is a binary variable equal to 1 when unit i is treated. Equation (1) illustrates a scenario where unit i 's outcome is not only influenced by its own treatment, represented by β_{it} , but also by the treatments of other units, as captured by γ_{it}^j and η_{it}^j . Assuming that the treatment effect in the absence of interference is homogeneous across units and time, we set $\beta_{it} = -0.5$ for all i and t , in which case the ATT without interference equals to -0.5 . We also assume that the time effect is given by $\delta_t = 0.1 \cdot (t - 1)$.

The left panel in Figure 2 illustrates a data pattern in scenarios without a spillover effect. Estimators that account for staggered adoption and heterogeneous treatment




Figure 1: Illustration of the potential treatment and spillover paths

effects typically utilize the observations from the never-treated group Z and the not-yet-treated observations in group B at time 2 as the control group. These observations are used to estimate the time trend and are contrasted with treated observations to estimate the ATT without interference. For example, [Wooldridge \[2022\]](#) proposes the following variant of the TWFE regression model. Let G_i be the group membership of unit i , with $G_a = G_{a'} = A$, $G_b = G_{b'} = B$, and $G_z = G_{z'} = Z$. The model is given by:

$$\begin{aligned}
 Y_{it} = & \alpha_i + \delta_t + \beta_{A2} \cdot \mathbf{1}(G_i = A, t = 2) \\
 & + \beta_{A3} \cdot \mathbf{1}(G_i = A, t = 3) \\
 & + \beta_{B3} \cdot \mathbf{1}(G_i = B, t = 3) + \varepsilon_{it},
 \end{aligned} \tag{2}$$

where, abusing notation, $(\beta_{A2}, \beta_{A3}, \beta_{B3})$ are coefficients on group-period indicators, and α_i, δ_t are the unit and time fixed effects, respectively. Under suitable conditions, it can be shown that the estimate of β_{gt} , denoted by $\hat{\beta}_{gt}$, is consistent for the ATT for each group g at each time t , all equal to -0.5 in our example.³

We now introduce spillovers under two alternative scenarios. We will also use these so-

³In fact, since the treatment is homogeneous, the standard TWFE regression would also consistently estimate the ATT.




Figure 2: Possible data patterns under Equation (1)

called Examples 1 and 2 throughout the paper to motivate our assumptions and empirical application. In the first scenario, the spillover is in the form of a diffusion effect, meaning that the direct effect β_{it} and the spillover effect $(\gamma_{it}^j, \eta_{it}^j)$ have the same sign. In the second scenario, the spillover is in the form of displacement, where β_{it} and $(\gamma_{it}^j, \eta_{it}^j)$ have opposite signs.

Example 1 (installation of a water treatment plant). Consider a scenario where we are interested in the effect of introducing a water treatment plant on the health outcomes of villages situated along a river. Suppose the nearby villages a and a' , categorized as group A, are the first to adopt the plant. This adoption not only improves water quality in these villages but may also enhance the water quality of the not-yet-treated downstream villages, resulting in a spillover effect.

The middle panel in Figure 2 visualizes a possible data pattern of Example 1. For a numerical illustration, let's build upon Equation (1) and set the spillovers to also be homogeneous across units and time: $\gamma_{it}^j = \eta_{it}^j = \gamma = \eta = -0.05$ for all i, j and t . There are two key challenges arising from this setting. First, there is no valid control group because both never-treated and not-yet-treated observations are negatively affected by spillovers. To illustrate, note that the time-difference of Equation (1) for an untreated unit is given by:

$$Y_{it} - Y_{i,t-1} = \delta_t - \delta_{t-1} + \sum_{j \neq i} (\gamma_{it}^j \cdot D_{jt} - \gamma_{i,t-1}^j \cdot D_{j,t-1}).$$

This equation reveals that the spillover effect introduces a bias term, disrupting the consistent estimation of the time trend $(\delta_t - \delta_{t-1})$. For instance, the bias term for the time-difference of an untreated unit z between periods 2 and 3 is calculated as $\gamma_{z3}^a + \gamma_{z3}^{a'} + \gamma_{z3}^b + \gamma_{z3}^{b'} - \gamma_{z2}^a - \gamma_{z2}^{a'} = -0.1$. Unit z' has the same bias term.

Second, even if we could correctly identify time and group fixed effects, it would not be possible to separately identify the direct and spillover effects. Estimators that account for staggered adoption and heterogeneous treatment effects, such as (2), would at best identify the average sum of the direct and spillover effects. For example, such an approach would estimate $\widehat{\beta}_{A2} = (\beta + \gamma) = -0.55$ and $\widehat{\beta}_{A3} = \widehat{\beta}_{B3} = (\beta + 3 \times \gamma) = -0.65$.

Example 2 (installation of stolen vehicle recovery devices). [Gonzalez-Navarro \[2013\]](#) studied the effect of installing a stolen vehicle recovery device on car theft incidents. The introduction of this treatment was staggered across different states within a country and was limited to specific car models. In this scenario, car theft could potentially be displaced to other unprotected models within treated states or to the same models in states that had not yet adopted the device. [Gonzalez-Navarro \[2013\]](#) found a 52% increase in thefts for the same models in states without the installed device.

The right panel in Figure 2 visualizes a possible data pattern of Example 2, where we set $\gamma_{it}^j = \eta_{it}^j = 0.05$ for all i, j and t . Example 2 face the same key challenges that we discussed earlier: the absence of a valid control group and the difficulty in separately estimating direct effects and spillover effects. Note that, especially in the case of displacement, spillover effects could intensify over time as more and more treated units spill on an increasingly narrower pool of untreated units.

While the sum of direct and spillover effects might be of interest in some cases, identifying the direct effect separately should be of prime importance in most contexts. For example, when a unit decides whether to participate in a policy or treatment, its main concern probably is the direct effect, because other units' decisions are out of its control. Policymakers whose jurisdiction spans all units might also want to understand the distinct impact of each channel. In addition, it should be noted that the sum of direct and spillover effects might have limited external validity, as this sum is specific to the observed treatment histories of all units, while there is a vast array of counterfactual treatment histories that all of these units might experience.

Figure 3 visualizes our key assumptions that allow the identification of the direct effect. They assume that all treated units are not influenced by spillovers, and that a subset of never-treated units remains unaffected by spillovers as well. Consequently, in this figure, there are no longer lines to treated observations, and there are no lines to unit z' , allowing for the identification of the time trend. In what follows, we detail how these




Figure 3: DAG under the key identification assumptions

assumptions are likely to hold in empirical applications, illustrated through Examples 1 and 2.

Example 1 [continued]. Consider villages situated at the most upstream part of a river, none of which have water treatment plants. These upstream villages are not affected by the installation of water treatment plants in other villages along the river, since all other villages are downstream relative to them. Therefore, in this context, these upstream villages represent untreated units that are not subject to spillover effects.

Next, consider the village located furthest downstream, which initially does not have a water treatment plant. When an upstream village installs a plant, the downstream village experiences spillover effects, benefiting from improved water quality resulting from the upstream water treatment. However, once the downstream village installs its own water treatment plant, the treatment status of the upstream village becomes irrelevant. The water quality in the downstream village is now only determined by its own treatment. Consequently, in this situation, treated units do not experience spillover effects.

Example 2 [continued]. Consider states that are distant from all states where stolen vehicle recovery devices have been installed in specific car models. These states might be unaffected by spillover effects, because car thieves deterred from targeting models

equipped with the device are likely to limit their alternative targets to those in areas within a manageable distance, for instance, because their networks are more robust. [Gonzalez-Navarro \[2013\]](#) shows that the data supports the notion that geographical constraints limit displacement behavior.

Next, consider a car model without the device, located in a state adjacent to the one where the device had been installed. This car model is subject to spillover effects because installing the device in the neighbouring state prompts thieves to redirect their targets to models without the device in nearby areas. However, once the device is installed in these previously unprotected models, they no longer experience spillover effects, as thieves' attention turns to vehicles still lacking the device.

It might be argued that as the coverage of states and car models with the protection device expands to become almost universal, thieves might eventually revert to targeting protected cars, violating the assumption. This scenario might not be totally implausible unless thieves shift their focus to other, less protected assets or leave the criminal market entirely. Nevertheless, such an almost universal adoption of the protection device would be considered an extreme case and hard to evaluate due to a very small set of control units.

3 Setup

We consider a DiD model with staggered treatment adoption, which involves panel data of units observed over time periods $t \in \{1, \dots, T\}$. For each unit at each time t , we consider a binary treatment status indicating whether the unit is treated (1) or not treated (0). We assume that the treatment is irreversible, meaning that once a unit undergoes treatment, it remains treated in all subsequent periods.

Assumption 1 (irreversibility). *For any two time periods (s, t) such that $s < t$, if a unit has a treatment status of 1 at time s , then it also has a treatment status of 1 at time t .*

Under Assumption 1, we can categorize units into groups according to the periods at which they enter treatment. Let \mathcal{G} be a subset of $\{1, \dots, T, \infty\}$ that represents the periods at which units enter treatment. A unit is assigned to group $g \in \mathcal{G}$ if it enters treatment at period g , except for the group labeled ∞ , which consists of units that remain untreated until time T .

We consider a population of units, indexed by i , for each group $g \in \mathcal{G}$. We denote unit i in group g by a (i, g) pair, and we let Λ_g represent the set of all (i, g) indices within group g in the population, with $\Lambda \equiv \bigcup_{g \in \mathcal{G}} \Lambda_g$ being the set of all indices across all groups. For each unit (i, g) , we define $D_{igt} \in \{0, 1\}$ as the binary treatment indicator

at time t , and we let $D_{ig} \equiv (D_{ig1}, \dots, D_{igT})$ represent the treatment history of this unit. Furthermore, we define the vector d_g as follows:

$$d_g \equiv \underbrace{(0, \dots, 0)}_{t < g}, \underbrace{(1, \dots, 1)}_{t \geq g}, \quad (3)$$

which represents the realized treatment history of the units in group g . We let d_g^t denote its treatment history up to time t . In cases with no ambiguity, we will use 0 to represent d_∞ , the realized treatment history of those who remain untreated until time T , since d_∞ corresponds to the vector of zeros.

Let $Y_{igt}(\{d_{jh}\}_{(j,h) \in \Lambda})$ be the potential outcome for unit (i, g) at time t when $\{D_{jh}\}_{(j,h) \in \Lambda}$ is set to $\{d_{jh}\}_{(j,h) \in \Lambda}$. It is important to note that the potential outcome depends on the treatment histories of all units in the population (Λ), whereas under SUTVA it would be a function of the unit's own treatment history only, i.e., $Y_{igt}(\{d_{jh}\}_{(j,h) \in \Lambda}) = Y_{igt}(d_{ig})$. To facilitate future discussions, we rewrite the potential outcome by partitioning the population's treatment into the unit's own treatment and those of the other units:

$$Y_{igt}(d_{ig}, \{d_{jh}\}_{(j,h) \in \Lambda \setminus \{(i,g)\}}).$$

This notation emphasizes the possibility of unit (i, g) being affected by spillover effects from units not included in the sample, as Λ represents the index set of the entire population. Note that Assumption 1 and the definition of the group labels \mathcal{G} imply that we observe $d_{jh} = d_h$ for every $(j, h) \in \Lambda$ in the data.

We define $\mathbf{d}_{(i,g)}$ to be a value of $\{d_{jh}\}_{(j,h) \in \Lambda \setminus \{(i,g)\}}$ where $d_{jh} = d_h$ for every $(j, h) \in \Lambda \setminus \{(i, g)\}$, representing the treatment histories for units other than (i, g) according to their group labels. In addition, we define $\mathbf{0}_{(i,g)}$ to be another value of $\{d_{jh}\}_{(j,h) \in \Lambda \setminus \{(i,g)\}}$ where $d_{jh} = 0$ for every $(j, h) \in \Lambda \setminus \{(i, g)\}$, representing absence of treatment for all units other than (i, g) .

These definitions lead to the following four types of potential outcomes that are relevant to our discussion:

- $Y_{igt}(d_g, \mathbf{d}_{(i,g)})$ represents the *observed* outcome where unit (i, g) is treated according to d_g , and other units (j, h) are treated according to $\mathbf{d}_{(i,g)}$.
- $Y_{igt}(d_g, \mathbf{0}_{(i,g)})$ represents the counterfactual outcome where unit (i, g) is treated according to the observed d_g , but all the other units (j, h) are untreated.
- $Y_{igt}(0, \mathbf{d}_{(i,g)})$ represents the counterfactual outcome where unit (i, g) is untreated, but other units (j, h) are treated according to the observed $\mathbf{d}_{(i,g)}$.

- $Y_{igt}(0, \mathbf{0}_{(i,g)})$ represents the counterfactual outcome where both unit (i, g) and all the other units (j, h) are untreated.

We assume that there is no anticipatory effect for these four types of potential outcomes, a standard assumption in DiD analyses.

Assumption 2 (no anticipation). $Y_{igt}(d_{ig}, \mathbf{d}_{(i,g)}) = Y_{igt}(d_{ig}^t, \mathbf{d}_{(i,g)}^t)$ where $d_{ig} \in \{0, d_g\}$ and $\mathbf{d}_{(i,g)} \in \{\mathbf{0}_{(i,g)}, \mathbf{d}_{(i,g)}\}$.

Under this assumption, we can refer to the not-yet-treated group, labelled as ∞ , as the never-treated group. Note that Assumption 2 still allows the treatment effect to be heterogeneous based on the duration of treatment exposure.

Next, we introduce the parallel trend assumption, specifically for a linear DiD model. We will extend our discussion to a nonlinear DiD model in a later section.

Assumption 3 (parallel trend, linear model). *For every group g at time t ,*

$$\mathbb{E}(Y_{igt}(\mathbf{0}^t, \mathbf{0}_{(i,g)}^t) | \alpha_{ig}) = \alpha_{ig} + \delta_t,$$

where $\alpha_{ig} \in \mathbb{R}$ is the unit fixed effect and $\delta_t \in \mathbb{R}$ is the common time effect such that $\delta_1 = 0$.

Assumption 3 can also be expressed in a standard form commonly found in the literature on DiD models (see, e.g., [Borusyak et al., 2021](#)):

$$Y_{igt}(\mathbf{0}^t, \mathbf{0}_{(i,g)}^t) = \alpha_{ig} + \delta_t + \varepsilon_{igt}, \quad (4)$$

where $\mathbb{E}(\varepsilon_{igt} | \alpha_{ig}) = 0$ for every group g at time t .

Without SUTVA, multiple definitions of the ATT arise. We first introduce the ATT without interference:

$$ATT_0(g, t) \equiv \mathbb{E}(Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(\mathbf{0}^t, \mathbf{0}_{(i,g)}^t)).$$

This definition of $ATT_0(g, t)$ captures the expected treatment effect at time t when unit (i, g) is the only treated unit in the population, thereby excluding any spillover effects from the other units. In other words, $ATT_0(g, t)$ captures the direct effect from the treatment, illustrated by the solid edges in Figure 1. This aligns with the conventional definition of the ATT under SUTVA and is the estimand of interest in our paper. We can then define an aggregate ATT by $ATT_0 = \sum_{g,t} w_{gt} ATT_0(g, t)$, where w_{gt} is a weight chosen by the econometrician (see, e.g., [Callaway and Sant'Anna \[2020\]](#)).⁴

⁴Note that $ATT_0(g, t)$ is typically defined only for pairs (g, t) satisfying $t \geq g$. In this paper, we

We can also consider an alternative definition of the ATT that includes spillover effects:

$$ATT_S(g, t) \equiv \mathbb{E}(Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)).$$

This definition differs from $ATT_0(g, t)$ in that it incorporates the spillover effects from other treated units. Note that $ATT_S(g, t)$ includes the spillover effects from all units with group labels $g \leq t$.

We refer to the difference $ATT_S(g, t) - ATT_0(g, t)$ as the average spillover effect on the treated:

$$AST(g, t) \equiv \mathbb{E}(Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)).$$

Lastly, it is useful to define another estimand, which we refer to as the average spillover effect on the untreated:

$$ASUT(g, t) \equiv \mathbb{E}(Y_{it}(0^t, \mathbf{d}_{(i,g)}^t) - Y_{it}(0^t, \mathbf{0}_{(i,g)}^t)).$$

4 Identification

The discussion on the identification of $ATT_0(g, t)$ is structured into two steps. We first show that identifying $ATT_0(g, t)$ is equivalent to identifying the sum of the time effect and the spillover effect on the treated. The second step then introduces conditions that allow the identification of this sum. An implication of our assumptions is that it unifies the definitions of the ATT by implying that $ATT_0(g, t) = ATT_S(g, t)$.

We first present the necessary and sufficient condition for identifying $ATT_0(g, t)$ when spillovers are present.

Theorem 1. *Suppose that Assumptions 1 to 3 hold, and that all units are untreated at $t = 1$. Then, for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time $t \geq g$, the parameter $ATT_0(g, t)$ is identified if and only if $\delta_t + AST(g, t)$ is identified.*

Proof. Refer to the Appendix for the proof of this theorem and others that follow. \square

The proof of Theorem 1 shows that, for every (g, t) satisfying $t \geq g$:

$$\mathbb{E}(Y_{igt}) = \mathbb{E}(\alpha_{ig}) + \delta_t + ATT_0(g, t) + AST(g, t).$$

The intuition for Theorem 1 is that since $\mathbb{E}(\alpha_{ig})$ is identified from the data for group g at $t = 1$, it follows that identification of $ATT_0(g, t)$ requires knowledge of δ_t (the time effect)

extend its definition to also include pairs satisfying $t < g$, in which case $d_g^t = 0^t$, resulting in a trivial definition of $ATT_0(g, t) = 0$. We adopt this extension as it simplifies the notation in the proofs of our results.

and $AST(g, t)$ (the average spillover effect on the treated). In general, Assumptions 1 to 3 are not sufficient for the identification of these two parameters. Note that $AST(g, t) = 0$ in the absence of spillover effects, in which case the identification of the $ATT_0(g, t)$ only requires knowledge of the time effect.

In what follows, we propose two additional assumptions that enable identification of $ATT_0(g, t)$. We state the first assumption below.

Assumption 4 (No spillover effects on treated units). *For every (g, t) such that $t \geq g$,*

$$Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) = Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t).$$

This assumption requires that once a unit receives treatment, it is no longer influenced by spillover effects. This means the unit forfeits any spillovers it may have previously received and remains unaffected by spillovers from subsequently treated groups. Recall that we have previously discussed the plausibility of this assumption in Section 2, illustrated through Examples 1 and 2. Note that Assumption 4 implies $AST(g, t) = 0$, and therefore $ATT_0(g, t) = ATT_S(g, t)$, unifying the definition of $ATT(g, t)$.

Next, we state the second assumption. For every group $g \in \mathcal{G}$, let $\Lambda_g^0 \subseteq \Lambda_g$ be a collection of units such that, for every untreated period $t < g$:

$$\mathbb{E}(Y_{igt}(0^t, \mathbf{d}_{(i,g)}^t) | \alpha_{ig}, (i, g) \in \Lambda_g^0) = \mathbb{E}(Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t) | \alpha_{ig}, (i, g) \in \Lambda_g^0) = \alpha_{ig} + \delta_t. \quad (5)$$

Hence, the Λ_g^0 set consists of units within group g that are not affected by spillover effects while they are untreated.

Assumption 5 (Existence of never-treated units without spillover effects). Λ_∞^0 has a positive measure.

This assumption states that there exists a nontrivial proportion of never-treated units that are not affected by spillovers, allowing for the identification of the time effect δ_t . In practice, the researcher may not have complete knowledge of Λ_g^0 and take a conservative approach by selecting the smaller subset of units strongly believed to be unaffected by spillovers, denoted by $\tilde{\Lambda}_g^0 \subseteq \Lambda_g^0$. For brevity of notation, we use Λ_g^0 interchangeably with $\tilde{\Lambda}_g^0$. All the results we discuss below apply to both Λ_g^0 and $\tilde{\Lambda}_g^0$.

Note that Assumption 5 does not impose any requirement about the size of Λ_g^0 for $g \neq \infty$. For instance, in the study by [Gonzalez-Navarro \[2013\]](#) described in Example 2, the author used the never-treated states that are farthest from the treated ones as controls, implying $\Lambda_g^0 = \emptyset$ for $g \neq \infty$. Alternatively, assuming that spillover effects occur only among adjacent states, Λ_g^0 could consist of all untreated observations in group g that are not adjacent to any treated state. In this case, the ‘‘control group’’ for time t , defined

by $\bigcup_{g>t} \Lambda_g^0$, decreases in t as more states adopt the treatment over time, resulting in a smaller number of untreated states that are not adjacent to any treated ones.

We conclude this section by showing that $ATT_0(g, t)$ is identified under these two additional assumptions, which is a direct consequence of Theorem 1.

Theorem 2. *Suppose that Assumptions 1 to 5 hold, and that all units are untreated at $t = 1$. Then, $ATT_0(g, t)$ is identified for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time $t \geq g$.*

5 Estimation and Inference

In this section, we discuss estimation and inference of $ATT_0(g, t)$ under Assumptions 1 to 5. Consider a balanced panel of T periods, where all units are untreated at $t = 1$. The units are indexed as $i = 1, \dots, N_g$ for each group label $g \in \mathcal{G}$.

For estimation, it is useful to introduce a binary variable S_{igt} that indicates whether an observation (i, g, t) could be subject to spillover effects. To define this variable, note that an observation is potentially influenced by spillovers under the following conditions:

- The observation is within the post-treatment period ($t \geq \min\{t \mid t \in \mathcal{G}\}$).
- The observation is not treated (with $D_{igt} = 0$), as otherwise treated observations are not influenced by spillover effects by Assumption 4.
- The observation does not belong to the set $\{\Lambda_g^0\}_{g \in \mathcal{G}}$, as otherwise units in these sets are considered unaffected by spillover effects, as defined in Equation (5).

Considering these, and defining $q \equiv \min\{t \mid t \in \mathcal{G}\}$ as the initial post-treatment period, we define S_{igt} as

$$S_{igt} = \begin{cases} 1 & \text{if } t \geq q \text{ and } D_{igt} = 0 \text{ and } (i, g) \notin \Lambda_g^0 \\ 0 & \text{otherwise} \end{cases}.$$

We first discuss estimation of $ATT_0(g, t)$ in the case where Λ_∞^0 is the only nonempty set. We propose the following extension of Wooldridge [2022] as the estimation procedure. Consider a partition of group ∞ into subgroups $(\infty, 0)$ and $(\infty, 1)$, where $(\infty, 0)$ consists of units belonging to Λ_∞^0 , and $(\infty, 1)$ consists of units belonging to $\Lambda_\infty - \Lambda_\infty^0$. With this partitioning, units are now categorized into a more extended set of groups than \mathcal{G} , denoted by $\tilde{\mathcal{G}}$. For example, if $\mathcal{G} = \{q, \dots, T, \infty\}$, then the extended set $\tilde{\mathcal{G}}$ is defined as

$$\tilde{\mathcal{G}} \equiv \{q, \dots, T, (\infty, 0), (\infty, 1)\}.$$

We use $\mathbf{g} \in \tilde{\mathcal{G}}$ to denote a generic group label within $\tilde{\mathcal{G}}$, and we extend the definitions of Y_{igt} , D_{igt} , and S_{igt} to this group label \mathbf{g} .

With these notations, we estimate the linear regression model where Y_{igt} is the outcome variable, and the regressors are:

- indicators of \mathbf{g} (the “extended group fixed effects”),
- indicators of t (the “time fixed effects”),
- interactions between D_{igt} and indicators of (\mathbf{g}, t) , and
- interactions between S_{igt} and indicators of (\mathbf{g}, t) .

In other words, we estimate the regression model:

$$\begin{aligned}
Y_{igt} = & \alpha_{\mathbf{g}} + \delta_t + \sum_{g' \in \mathcal{G} \setminus \{\infty\}} \sum_{t'=g'}^T \beta_{g't'} \cdot \mathbf{1}((\mathbf{g}, t) = (g', t')) \cdot D_{igt} \\
& + \sum_{\mathbf{g}' \in \tilde{\mathcal{G}} \setminus \{(\infty, 0)\}} \sum_{t'=q}^{\mathbf{g}'-1} \gamma_{\mathbf{g}'t'} \cdot \mathbf{1}((\mathbf{g}, t) = (\mathbf{g}', t')) \cdot S_{igt} + \varepsilon_{igt},
\end{aligned} \tag{6}$$

where $\mathbf{g} - 1$ is interpreted as T if $\mathbf{g} = (\infty, 1)$. Then, $\hat{\beta}_{gt}$ is the estimate of $ATT_0(g, t)$.⁵

Note that Equation (6) involves $\alpha_{\mathbf{g}}$, the *group* fixed effect, as opposed to α_{ig} , the *unit* fixed effect. This applies similarly to the treatment effects, where Equation (6) involves group-level treatment effects (β_{gt}) instead of unit-level treatment effects. This simplifies the estimation and inference of $ATT_0(g, t)$, because the estimate $\hat{\beta}_{gt}$ and its standard error can be easily obtained through standard linear regression in any statistical software package. Moreover, estimation and inference of an aggregate ATT is also straightforward, because the estimate is given by $\sum_{g,t} w_{gt} \hat{\beta}_{gt}$, and its standard error is straightforwardly computed by

$$\text{Var} \left(\sum_{g,t} w_{gt} \hat{\beta}_{gt} \right) = \sum_{g,t} \sum_{g',t'} w_{gt} w_{g't'} \text{Cov}(\hat{\beta}_{gt}, \hat{\beta}_{g't'}),$$

where $\text{Cov}(\hat{\beta}_{gt}, \hat{\beta}_{g't'})$ is available in any statistical software package, e.g., via $\mathbf{e}(\mathbf{V})$ in Stata.

Alternatively, the following extension of the imputation-based procedure of [Borusyak et al. \[2021\]](#) offers a numerically equivalent method of obtaining $\hat{\beta}_{gt}$.

1. Estimate the linear model

$$Y_{igt} = \alpha_{ig} + \delta_t + \varepsilon_{igt},$$

⁵In practice, when implementing Equation (6) using a statistical software package, the researcher may simply ignore the limits of the summation terms. Instead, one may include the interaction terms for all $\mathbf{g} \in \tilde{\mathcal{G}}$ and for every $1 \leq t \leq T$. The software will automatically omit the extra interactions from the model due to the multicollinearity.

using observations (i, g, t) such that $D_{igt} = 0$ and $S_{igt} = 0$.⁶

- Let $\widehat{\alpha}_{ig}$ and $\widehat{\delta}_t$ be the estimates of α_{ig} and δ_t from the linear model. Impute the baseline outcome for unit (i, g) at time t as

$$\widehat{Y}_{igt}(0^t, \mathbf{0}_{(i,g)}^t) = \widehat{\alpha}_{ig} + \widehat{\delta}_t.$$

- For a treated group g at time $t \geq g$, estimate $ATT_0(g, t)$ by

$$\frac{1}{N_g} \sum_{i=1}^{N_g} \left[Y_{igt} - \widehat{Y}_{igt}(0^t, \mathbf{0}_{(i,g)}^t) \right],$$

which can be shown to be equal to $\widehat{\beta}_{gt}$.

In this procedure, $ATT_0(g, t)$ is estimated by the average difference between the observed (treated) outcome and the counterfactual (baseline) outcome. The baseline outcome is estimated using observations in the control group that are unaffected by spillovers.

Note that the estimate of $ATT_0(g, t)$ in the imputation-based procedure equals to

$$\frac{1}{N_g} \sum_{i=1}^{N_g} Y_{igt} - \frac{1}{N_g} \sum_{i=1}^{N_g} \widehat{\alpha}_{ig} - \widehat{\delta}_t.$$

The regression in Equation (6) directly calculates $(1/N_g) \sum_{i=1}^{N_g} \widehat{\alpha}_{ig}$, rather than individual $\widehat{\alpha}_{ig}$ values, through the group-level fixed effect. The following result shows that, despite this simplification, the population regression of Equation (6) correctly identifies $ATT_0(g, t)$. The consistency and asymptotic normality of $\widehat{\beta}_{gt}$ follows directly from the validity of the population regression.

Theorem 3. *Suppose that the assumptions of Theorem 2 hold. Consider the population regression of Equation (6), and let β_{gt} be the population regression coefficient for the interaction between D_{igt} and the indicator of (g, t) . Then, $\beta_{gt} = ATT_0(g, t)$.*

Next, we consider the case where Λ_g^0 might be non-empty for some $g \neq \infty$. We define the extended set of groups $\tilde{\mathcal{G}}$ by

$$\tilde{\mathcal{G}} \subseteq \mathcal{G} \times \{0, 1\}.$$

This partitions each group g into subgroups $(g, 0)$ and $(g, 1)$ whenever Λ_g^0 is nonempty, where $(g, 0)$ consists of units belonging to Λ_g^0 and $(g, 1)$ consists of units belonging to

⁶In balanced panels, it is sufficient to use the group fixed effect α_g in the linear model.

$\Lambda_g - \Lambda_g^0$. We then extend the definition of Y_{igt} , D_{igt} , and S_{igt} to the group label $\mathbf{g} \in \tilde{\mathcal{G}}$ and define ATT_0 accordingly:

$$ATT_0(\mathbf{g}, t) = \mathbb{E}(Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)).$$

The aggregate ATT can then be defined as $ATT_0 = \sum_{\mathbf{g}, t} w_{\mathbf{g}t} ATT_0(\mathbf{g}, t)$, where $w_{\mathbf{g}t}$ is a weight chosen by the econometrician. The regression in (6) can be straightforwardly extended to this group label, with the modification that the coefficient $\beta_{\mathbf{g}t}$ now represents $\beta_{\mathbf{g}t} = ATT_0(\mathbf{g}, t)$.

Lastly, if the data is an unbalanced panel, the regression in Equation (6) is no longer consistent for the ATT_0 . The imputation-based estimation procedure of [Borusyak et al. \[2021\]](#) is still consistent, but the standard error will be asymptotically conservative in general (see [Borusyak et al., 2021](#), Section 4.3). In contrast, for a balanced panel, the standard error computed from Equation (6) is asymptotically exact.

6 Extension to Nonlinear DiD Models

In this section, we extend our previous findings to the case where Y_{igt} is a count variable, for which the linear parallel trend condition (Assumption 3) does not hold. This extension contributes to the literature on nonlinear DiD models [[Wooldridge, 2023](#)], expanding the applicability of our results to a wider array of empirical applications.

We introduce the following assumption regarding parallel trends in the context of count data.

Assumption 3' (parallel trend, Poisson model). *For every group g at time t ,*

$$\ln \mathbb{E}(Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t) | \alpha_{ig}) = \alpha_{ig} + \delta_t.$$

By replicating the arguments in Theorems 1 and 2, the following corollaries show that $ATT_0(g, t)$ is identified under assumptions similar to those in Theorem 2. In doing so, we abuse notation and define $\Lambda_g^0 \subseteq \Lambda_g$ for every group $g \in \mathcal{G}$ as a collection of units such that, for every untreated period $t < g$:

$$\ln \mathbb{E}(Y_{igt}(0^t, \mathbf{d}_{(i,g)}^t) | \alpha_{ig}, (i, g) \in \Lambda_g^0) = \ln \mathbb{E}(Y_{igt}(0^t, \mathbf{0}^t) | \alpha_{ig}, (i, g) \in \Lambda_g^0) = \alpha_{ig} + \delta_t. \quad (7)$$

Corollary 1. *Suppose that Assumptions 1 and 2 and assumption 3' hold, and that all units are untreated at $t = 1$. Then, for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time $t \geq g$, the $ATT_0(g, t)$ is identified if and only if $\mathbb{E}(\exp\{\alpha_{ig}\}) \cdot \exp\{\delta_t\} + AST(g, t)$ is identified.*

Corollary 2. *Suppose that Assumptions 1, 2, 4 and 5 and assumption 3' hold, and that all units are untreated at $t = 1$. Then, $ATT_0(g, t)$ is identified for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time $t \geq g$.*

Note that, despite a nonlinear setting, the identification holds in a short panel setting where T remains fixed.

Let S_{igt} be defined as in previous sections, and consider a balanced panel of T periods where units are indexed as $i = 1, \dots, N_g$ for each group label g , and all units are untreated at $t = 1$. Our parameter of interest is still $ATT_0(g, t)$. In the case of count data, the average treatment effect in terms of percentage changes is also often reported:

$$ATTP_0(g, t) = \frac{ATT_0(g, t)}{\mathbb{E}(Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t))},$$

which can be aggregated to define an $ATTP_0 \equiv \sum_{g,t} w_{gt} ATTP_0(g, t)$.

The estimation and inference procedure discussed in Section 5 can be straightforwardly extended to the count data. For example, in the case where Λ_∞^0 is the only nonempty set, we define the extended group label \mathbf{g} to be as defined in Section 5, and we use the following simple estimation procedure that involves a parsimonious generalized linear model.

1. Estimate the Poisson regression model where Y_{igt} is the outcome variable, and the regressors are:
 - indicators of \mathbf{g} (the “extended group fixed effects”),
 - indicators of t (the “time fixed effects”),
 - interactions between D_{igt} and indicators of (\mathbf{g}, t) , and
 - interactions between S_{igt} and indicators of (\mathbf{g}, t) .

In other words, we estimate the Poisson regression model:

$$\begin{aligned} \ln \mathbb{E}(Y_{igt} | \mathbf{X}_{igt}) = & \alpha_{\mathbf{g}} + \delta_t + \sum_{g' \in \mathcal{G} \setminus \{\infty\}} \sum_{t'=g'}^T \beta_{g't'} \cdot \mathbf{1}((\mathbf{g}, t) = (g', t')) \cdot D_{igt} \\ & + \sum_{\mathbf{g}' \in \tilde{\mathcal{G}} \setminus \{(\infty, 0)\}} \sum_{t'=q}^{\mathbf{g}'-1} \gamma_{\mathbf{g}'t'} \cdot \mathbf{1}((\mathbf{g}, t) = (\mathbf{g}', t')) \cdot S_{igt}, \end{aligned} \quad (8)$$

where $\mathbf{g} - 1$ is interpreted as T if $\mathbf{g} = (\infty, 1)$ and \mathbf{X}_{igt} represents the vector of regressors. Let $\hat{\alpha}_{\mathbf{g}}$, $\hat{\delta}_t$, and $\hat{\beta}_{gt}$ be the estimates of $\alpha_{\mathbf{g}}$, δ_t , and β_{gt} from this model, respectively.

2. Estimate $ATT_0(g, t)$ by

$$\widehat{ATT}_0(g, t) = \exp\{\widehat{\alpha}_g + \widehat{\delta}_t + \widehat{\beta}_{gt}\} - \exp\{\widehat{\alpha}_g + \widehat{\delta}_t\},$$

or estimate $ATTP_0(g, t)$ by $\widehat{ATTP}_0(g, t) = \exp\{\widehat{\beta}_{gt}\} - 1$.

The validity of the population regression of Equation (8) can be shown by replicating the arguments in Theorem 3, and we omit the proof here. The consistency and asymptotic normality of $\widehat{ATT}_0(g, t)$ and $\widehat{ATTP}_0(g, t)$ follow directly from the validity of the population regression.

Note that most statistical software packages that run Poisson regressions calculate the standard errors of $(\widehat{\alpha}_{\mathbf{g}}, \widehat{\delta}_t, \widehat{\beta}_{gt})$ using the maximum likelihood. This assumes that the distribution of $Y_{igt}(0^t, \mathbf{0}_{(i,\mathbf{g})}^t)$ conditional on $\alpha_{\mathbf{ig}}$ follows a Poisson distribution (as opposed to only specifying its mean as in Assumption 3'), ruling out heteroskedasticity. To accommodate heteroskedasticity, standard errors can instead be derived using the quasi-maximum likelihood estimation (QMLE) method. Specifically, let θ be the vector of all coefficients in the Poisson regression (i.e., all of α_g , δ_t , β_{gt} , and κ_{gt}), $\hat{\theta}$ be their maximum likelihood estimates (i.e., all of $\widehat{\alpha}_{\mathbf{g}}$, $\widehat{\delta}_t$, $\widehat{\beta}_{gt}$, and $\widehat{\gamma}_{gt}$), and \mathbf{X}_{igt} be the vector of all regressors. Let $\{\Lambda^c\}_{c=1}^C$ be the partition of units according to which the units are clustered. Define

$$\mathcal{S} = \sum_{c=1}^C \left[\sum_{(i,\mathbf{g}) \in \Lambda^c} \sum_{t=1}^T \mathbf{X}_{igt} (Y_{igt} - \widehat{Y}_{igt}) \right] \left[\sum_{(i,\mathbf{g}) \in \Lambda^c} \sum_{t=1}^T \mathbf{X}_{igt} (Y_{igt} - \widehat{Y}_{igt}) \right]'$$

as the clustered outer product of the score function, where $\widehat{Y}_{igt} = \exp\{X'_{igt}\hat{\theta}\}$ is the fitted value of Y_{igt} in the Poisson regression.⁷ In addition, define

$$\mathcal{H} = \sum_{c=1}^C \sum_{(i,\mathbf{g}) \in \Lambda^c} \sum_{t=1}^T \mathbf{X}_{igt} \mathbf{X}'_{igt} \widehat{Y}_{igt}$$

as the negative Hessian function. Then, the variance-covariance matrix of $\hat{\theta}$ is given by

$$\widehat{\text{Var}}(\hat{\theta}) = \mathcal{H}^{-1} \mathcal{S} \mathcal{H}^{-1}.$$

This variance-covariance matrix can then be used to compute the standard errors of the ATT_0 and $ATTP_0$ estimates via the delta method.

⁷We abuse notation and let \widehat{Y}_{it} represent a different object from the linear case.

7 Application to Auto Theft Prevention Policy

In this section, we apply our method to revisit the findings of [Gonzalez-Navarro \[2013\]](#), who studied the effects of installing an auto theft prevention device known as Lojack. This was a compact device installed in vehicles, allowing for tracking of the vehicle.

The policy was implemented in Mexico through an exclusive agreement between the Ford Motor Company and the Lojack company. Initially, the technology was introduced for a particular Ford car model (Ford Windstar) in a specific state (Jalisco) among the 2001 car models. Subsequently, the installation of Lojack expanded to include other $model \times state$ combinations, eventually encompassing 32 $model \times state$ combinations by 2004. The dataset of [Gonzalez-Navarro \[2013\]](#) provides comprehensive information on car theft for each $model \times state \times vintage$ (the car model’s year) combination, for each calendar year. For our analysis, we use the indices m , s , v , and t to represent car model, state, vintage, and the calendar year of the auto theft, respectively.

[Gonzalez-Navarro \[2013\]](#) points out two possible sources of spillover effects following the introduction of Lojack. The first potential source is within-state spillover to car models not equipped with Lojack. Given the public knowledge about specific car models and states where Lojack was installed, criminals may alter their target preferences, focusing on car models without Lojack within the same state. The second source is geographical spillovers, where installing Lojack in certain models may prompt thieves, particularly those specializing in those models, to shift their operations to other states where these specific models remain unprotected by Lojack.

Because of the potential for such spillovers, [Gonzalez-Navarro \[2013\]](#) relies only on time-series variation for identification, illustrating the challenge in extending the DiD framework to spillovers:

“In the presence of spatial externalities, DiD estimation using observations from different geographical locations produces biased estimates of policy impact. The basic challenge is that whenever treatment in one geographical location also has effects in control locations, these are no longer valid counterfactual observations. Furthermore, DiD estimation precludes actual estimation of externalities unless there is a set of observations subject to externalities and a set of observations that is not, so that the latter can play the role of counterfactual. For these reasons I do not use DiD estimation. Instead, I use an interrupted time series strategy in which the counterfactual is given by observations occurring before the intervention.”

Nevertheless, as a robustness check, [Gonzalez-Navarro \[2013\]](#) also estimates a DiD model while attempting to control for spillover effects, but without accounting for the

staggered adoption design. In this section, we apply our method to revisit this study and estimate the treatment effect across various combinations of groups and time periods, thereby revealing the heterogeneous effects of Lojack installation.

Once Lojack was installed in a particular combination of car model, state, and vintage, it continued to be installed in all subsequent vintages of that model in the same state. This setup allows us to treat the situation as a staggered adoption design, where the unit of analysis is defined as the combination of *model* (m) \times *state* (s) \times *age* (a). *age* refers to the number of years elapsed since the car’s model year, calculated as the difference between the calendar year (t) and the vintage year (v), such that $a = t - v$. Under this framework, our analysis is based on a balanced panel subset derived from the original dataset, consisting of 1152 units observed over 6 years from 1999 to 2004.

We define the binary treatment indicator for a unit (m, s, a) at time t as D_{msat} . To illustrate, consider the Ford Windstar model in Jalisco. For this unit, Lojack has been installed in all newly released ($age = 0$) vehicles starting in 2001. Thus, for a Ford Windstar model in Jalisco with $age = 0$, we have $D_{Windstar, Jalisco, 0, t} = 1$ for every $t \geq 2001$.

Our method relies on Assumptions 4 and 5. Assumption 4 requires that once a *model* \times *state* \times *age* unit has Lojack installed, it is not influenced by spillover effects. Generally, when Lojack is installed in certain units, we can expect that thieves targeting those models will shift their focus towards vehicles without Lojack protection, rather than those already with Lojack. Thus, it is reasonable to assume that units already fitted with Lojack will not be subject to displacement effects from other units, satisfying Assumption 4. Assumption 5 requires that there exist units which are not affected by spillover effects, and Gonzalez-Navarro [2013] provides empirical support for this assumption, demonstrating that car models in states geographically distant from those where the treatment was applied do not experience spillover effects.⁸

Let Y_{msat} be the number of auto thefts for a *model* \times *state* \times *age* unit that occurred in a given calendar year t . We consider two kinds of empirical models for this outcome. First, we consider a linear parallel trend:

$$\mathbb{E}(Y_{msat}(0^t, \mathbf{0}_{(msa,t)}^t) | \alpha_{msa}) = \alpha_{msa} + \delta_t.$$

This is equivalent to Assumption 3, where the combination (m, s, a) plays the role of

⁸The results of Gonzalez-Navarro [2013] using only time series variation vs. the DID approach are similar, suggesting that the units in states distant from the treated areas are unaffected by the installation of Lojack.

(i, g). Second, we consider a Poisson parallel trend:

$$\ln \mathbb{E}(Y_{msat}(0^t, \mathbf{0}_{(msa,t)}^t) | \alpha_{msa}) = \alpha_{msa} + \delta_t,$$

which is equivalent to Assumption 3'. The second model is particularly suitable when Y_{msat} is a count variable with a high frequency of zeros, in which case a Poisson regression model is more appropriate.

We define the Λ_∞^0 set as the collection of (m, s, a) units where s is a state that is not adjacent to any state with treated units throughout the rollout of Lojack. We then define S_{msat} as a binary indicator that is equal to 1 if $t \geq 2001$, $D_{msat} = 0$ and $(m, s, a) \notin \Lambda_\infty^0$. In addition, define $\bar{\mathcal{G}} = \{2001, 2002, 2003, 2004\}$ as the set of group labels for treated units, categorized by the period at which units enter treatment. Let N_g be the number of units in group $g \in \bar{\mathcal{G}}$ within the dataset, and let $\bar{N} \equiv \sum_{g=2001}^{2004} \sum_{t=g}^{2004} N_g = \sum_{g=2001}^{2004} (2005 - g)N_g$ be the total number of treated observations in the dataset. We estimate the following aggregate ATT_0 s:

$$\begin{aligned} ATT_0 &= \sum_{g=2001}^{2004} \sum_{t=g}^{2004} \frac{N_g}{\bar{N}} ATT_0(g, t), \\ ATT_0^0 &= \sum_{g=2001}^{2004} \frac{N_g}{N_{2001} + \dots + N_{2004}} ATT_0(g, g), \\ ATT_0^1 &= \sum_{g=2001}^{2003} \frac{N_g}{N_{2001} + \dots + N_{2003}} ATT_0(g, g + 1), \\ ATT_0^2 &= \sum_{g=2001}^{2002} \frac{N_g}{N_{2001} + N_{2002}} ATT_0(g, g + 2). \end{aligned}$$

Here, ATT_0 measures the overall effect of Lojack installation, computed as the weighted average of all $ATT_0(g, t)$ values across g and t . The ATT_0^k values, on the other hand, represent the weighted average of ATT_0 for the k -th year after installation of Lojack, measuring the temporal effects. For example, ATT_0^0 represents the immediate effect in the same year as the Lojack installation, ATT_0^1 represents the effect one year post-installation, and so forth.

Table 1 presents the estimated ATT_0 values obtained from both linear and Poisson model specifications, with standard errors clustered at the unit level. The analysis reveals a notable average reduction in thefts of 60% for the linear model and 64% for the Poisson model, highlighting Lojack's substantial deterrent effect. Moreover, the results from both models indicate that the rate of theft reduction becomes more pronounced over time, where the effect becomes statistically significant starting one year after installation. This

	Linear			Poisson		
	Estimate	Std Error	Reduction	Estimate	Std Error	Reduction
ATT_0	-6.1017	2.8893	-60%	-5.6349	2.5086	-66%
ATT_0^0	-3.9455	2.9166	-38%	-3.8738	2.4503	-50%
ATT_0^1	-6.7536	2.9801	-77%	-6.2742	2.5453	-77%
ATT_0^2	-16.9622	2.9691	-79%	-13.4790	4.2276	-85%

Table 1: Estimates of the aggregate ATT_0 s. The standard errors are clustered at the *model* (m) \times *state* (s) \times *age* (a) level. The “Reduction” column stands for the reduction rate, which is calculated using the formula for computing $ATTP_0$.

highlights the increasing effectiveness of Lojack in preventing auto thefts over time.

For comparison, we also report the estimated ATT_0 s from two misspecified models. First, we consider the TWFE specification that incorporates spillover effects but overlooks the staggered adoption nature of the treatment. Second, we consider the specification of Wooldridge [2022] and Borusyak et al. [2021] that accounts for staggered adoption but does not include spillover effects. The results from these models are presented in Table 2. We find that the TWFE regression estimate closely aligns with the estimates presented in Table 1. However, the estimates that neglect spillover effects exhibit an upward bias relative to the correctly specified estimates in Table 1. This is what we would expect in the presence of displacement effects, where installing Lojack in a treated unit increases theft for units without Lojack.

	TWFE-Linear		WB-Linear	
	Estimate	Reduction	Estimate	Reduction
ATT_0	-7.8595	-69%	-7.8335	-72%
ATT_0^0	N/A		-5.6375	-58%
ATT_0^1	N/A		-8.4526	-82%
ATT_0^2	N/A		-19.1385	-88%
	TWFE-Poisson		WB-Poisson	
	Estimate	Reduction	Estimate	Reduction
ATT_0	-5.4990	-61%	-5.8514	-62%
ATT_0^0	N/A		-3.9569	-43%
ATT_0^1	N/A		-6.4894	-73%
ATT_0^2	N/A		-14.5736	-94%

Table 2: Estimates of the aggregate ATT_0 s using the TWFE specification (the “TWFE” columns), and the specification of Wooldridge [2022] and Borusyak et al. [2021] (the “WB” columns), for each of linear and Poisson specifications. The “Reduction” columns stand for the reduction rate, which is calculated using the formula for computing $ATTP_0$.

8 Monte Carlo Simulation

In this section, we study the finite sample properties of our estimator in a simulated dataset, highlighting the bias-variance trade-off of our approach. We consider a balanced panel dataset over T periods, with either a simultaneous or staggered adoption design, starting with a pre-treatment period of $t = 1$. We consider M units in each group $g \in \mathcal{G} \equiv \{2, \dots, T, (\infty, 0), (\infty, 1)\}$, meaning that we have a total of $N = (T + 1)M$ units in the dataset. In the absence of spillover effects, our estimator is less efficient than conventional estimators that rule out spillover effects. However, in the presence of spillovers, the conventional estimators become biased. Given this bias-variance trade-off, when the sample size is small, the improvement in bias may not sufficiently offset the loss in precision.

Specifically, we consider the following data generating process (DGP) that embeds Assumptions 1 to 5. Depending on the specification of the outcome model—linear or Poisson—we adapt the relevant assumption, replacing Assumption 3 with 3' as necessary. The DGP is given by:

$$\mathbb{E}(Y_{igt} | \alpha_{ig}) = F \left(\alpha_{ig} + \delta_t + \beta_{igt} D_{igt} + (1 - D_{igt}) \cdot \sum_{h \in \mathcal{G} \setminus \{g\}} \sum_{j=1}^M \gamma_{igt}^{(j,h)} \cdot D_{jht} \right),$$

where the function F is $F(x) = x$ for the linear model or $F(x) = \exp(x)$ for the Poisson model, and $\gamma_{igt}^{(j,h)}$ represents the spillover effect from unit (j, h) to unit (i, g) . We parametrize the DGP as follows.

- The M units in each group $g \in \{2, \dots, T, (\infty, 0), (\infty, 1)\}$ are homogeneous, implying that $\alpha_{ig} = \alpha_g$, $\beta_{igt} = \beta_{gt}$ and $\gamma_{igt}^{(j,h)} = \gamma_{gt}^h$.
- Unit fixed effects are set to $\alpha_i = 26 - g + 1$ for all groups except for $(\infty, 0)$ and $(\infty, 1)$, where $\alpha_{(\infty,0)} = \alpha_{(\infty,1)} = 26 - T + 1$. This reflects selection into treatment because the units with earlier treatment have larger unit fixed effects. In the case of the Poisson model, we instead set $\alpha_g = \log(26 - g + 1)$.
- Common time effects are set to $\delta_t = \bar{\alpha} \times 0.1 \times ((t - 1) + \sin(t))$, where $\bar{\alpha}$ is the average of the unit fixed effects across all groups. This specification involves a linear upward trend $(t - 1)$ and a period-specific fluctuation modeled through $\sin(\cdot)$.
- The treatment effect is set to $\beta_{gt} = 0.5\alpha_g/t$. This effect is heterogeneous across groups and time periods, but homogeneous within a group. The effect gradually diminishes over time, with β_{gt} decreasing in t for each group g . The immediate


effect β_{gg} is largest for group $g = 2$ with the highest α_g . This parametrizes sorting on gain since α_g also correlates with treatment timing.

- Spillover effects are set to $\eta_{gt}^h = -\rho \cdot \beta_{gt}/U_t$, representing displacement effects, where U_t is the number of untreated units at time t except for those in $(\infty, 0)$. That is, for each treated unit (i, g) , we consider a total spillover effect of $-\rho \cdot \beta_{gt}$, where $\rho \in [0, 1]$ denotes the spillover intensity. This total effect is then evenly spread among all untreated units excluding those in $(\infty, 0)$. As a result, each untreated unit receives a spillover effect of $-\rho \cdot \beta_{gt}/U_t$ from the treated unit (i, g) .


With this parametrization, Y_{igt} is generated with an independent additive error term $\epsilon_{igt} \sim N(0, \max(\alpha_g)/10)$ for the linear model, and according to Poisson distribution for the Poisson model. We then estimate the aggregate ATT_0 defined as in Section 7, namely $ATT_0 = (1/\bar{G}) \sum_{g=2}^T \sum_{t=g}^T ATT_0(g, t)$, where $\bar{G} \equiv T(T-1)/2$ is the total number of treated group-time pairs in the dataset. We compare the mean Absolute Bias and the Mean Squared Error (MSE) across the following estimators:

- ($\hat{\beta}_1$) The TWFE estimator, which neither accounts for staggered treatment adoption nor for spillovers.
- ($\hat{\beta}_2$) The extended TWFE estimator by [Wooldridge \[2022\]](#), which accounts for staggered treatment adoption but does not account for spillovers. This estimator is numerically equivalent to the imputation estimator by [Borusyak et al. \[2021\]](#).
- ($\hat{\beta}_3$) Our estimator, which accounts for both staggered treatment adoption and spillovers.

Figure 4a and Table 3 present results from the linear DGP. The Figure visually contrasts the MSE across the three estimators to illustrate their relative performance under different scenarios, while the Table details their MSE and Absolute Bias values. Note that, when $T = 2$, the TWFE and the [Wooldridge \[2022\]](#) estimators are equivalent since treatment is not staggered. Overall, the relative performances of the estimators depend on the degree of spillovers, staggered treatment, and the number of units in each group. Intuitively, due to its efficiency, the TWFE has the lowest MSE in scenarios with no or little spillovers and with very few observations. As the number of observations increases and spillovers remain small, the [Wooldridge \[2022\]](#) estimator becomes the best-performing one, adjusting for staggered treatment without substantial bias. However, in scenarios where spillovers are not negligible and the number of units is large, our estimator achieves the lowest MSE, often by a large margin. Our estimator also performs better as treatment becomes more staggered ($T = 8$), highlighting our estimator’s ability to accurately account for cumulative spillovers affecting the untreated units’ outcomes. Furthermore,



(a) Linear



(b) Poisson

Figure 4: Comparison of the MSEs. Cell background color indicates the best-performing estimator. The numbers in cells represent the MSE ratios $\frac{MSE_1}{MSE_3}$ and $\frac{MSE_2}{MSE_3}$ respectively. The subscripts refer to: (1) TWFE estimator, (2) Wooldridge [2022] estimator, and (3) our estimator.

Table 3: MSE and Absolute Bias values - Linear

ρ	T	M	ATT	$ Bias_1 $	$ Bias_2 $	$ Bias_3 $	MSE_1	MSE_2	MSE_3	
0.00	2	1	-6.500	3.589	3.589	4.235	19.642	19.642	27.519	
		3	-6.500	2.060	2.060	2.374	6.613	6.613	8.957	
		5	-6.500	1.615	1.615	1.873	3.991	3.991	5.335	
		10	-6.500	1.159	1.159	1.313	2.134	2.134	2.745	
	8	1	-2.297	0.898	1.038	2.336	1.244	1.693	8.651	
		3	-2.297	0.554	0.606	1.463	0.477	0.569	3.336	
		5	-2.297	0.463	0.473	1.051	0.334	0.350	1.789	
		10	-2.297	0.384	0.334	0.782	0.217	0.172	0.954	
	0.25	2	1	-6.500	3.807	3.807	4.321	22.129	22.129	28.292
			3	-6.500	2.153	2.153	2.391	7.274	7.274	8.856
5			-6.500	1.715	1.715	1.824	4.584	4.584	5.280	
10			-6.500	1.316	1.316	1.342	2.642	2.642	2.805	
8		1	-2.297	1.403	1.335	2.456	2.776	2.717	9.587	
		3	-2.297	1.396	1.186	1.509	2.314	1.876	3.561	
		5	-2.297	1.355	1.116	1.114	2.054	1.559	1.917	
		10	-2.297	1.342	1.069	0.783	1.928	1.324	0.999	
0.50		2	1	-6.500	3.870	3.870	4.012	23.616	23.616	25.838
			3	-6.500	2.565	2.565	2.395	10.048	10.048	9.274
	5		-6.500	2.139	2.139	1.877	6.991	6.991	5.448	
	10		-6.500	1.796	1.796	1.338	4.700	4.700	2.787	
	8	1	-2.297	2.403	2.284	2.465	6.923	6.688	9.499	
		3	-2.297	2.367	2.184	1.440	5.985	5.317	3.220	
		5	-2.297	2.374	2.178	1.087	5.858	5.073	1.847	
		10	-2.297	2.404	2.212	0.782	5.900	5.070	0.970	
	0.75	2	1	-6.500	4.138	4.138	4.162	26.753	26.753	28.210
			3	-6.500	2.905	2.905	2.386	12.336	12.336	9.081
5			-6.500	2.652	2.652	1.869	9.928	9.928	5.476	
10			-6.500	2.405	2.405	1.330	7.524	7.524	2.768	
8		1	-2.297	3.448	3.346	2.581	13.089	12.935	10.172	
		3	-2.297	3.452	3.352	1.371	12.300	11.811	2.899	
		5	-2.297	3.417	3.290	1.099	11.884	11.162	1.912	
		10	-2.297	3.424	3.308	0.766	11.839	11.106	0.915	
1.00		2	1	-6.500	4.373	4.373	4.148	29.442	29.442	26.621
			3	-6.500	3.568	3.568	2.355	17.556	17.556	8.570
	5		-6.500	3.370	3.370	1.906	15.018	15.018	5.723	
	10		-6.500	3.305	3.305	1.270	12.749	12.749	2.555	
	8	1	-2.297	4.473	4.419	2.453	21.073	21.098	9.491	
		3	-2.297	4.445	4.399	1.394	20.124	19.900	3.076	
		5	-2.297	4.446	4.389	1.134	19.983	19.587	2.051	
		10	-2.297	4.486	4.426	0.802	20.246	19.771	1.020	

Note. Results over 1000 repetitions. Subscript refers to: (1) TWFE estimator, (2) [Wooldridge \[2022\]](#) estimator, and (3) our estimator. The lowest value across estimators is in bold.

Table 4: MSE and Absolute Bias values - Poisson

ρ	T	M	ATT	$ Bias_1 $	$ Bias_2 $	$ Bias_3 $	MSE_1	MSE_2	MSE_3	
0.00	1	1	-26.780	9.701	9.701	11.905	153.431	153.431	239.845	
		2	-26.780	5.540	5.540	6.740	47.762	47.762	73.687	
		5	-26.780	4.245	4.245	5.032	29.219	29.219	40.018	
	2	10	-26.780	3.071	3.071	3.689	14.968	14.968	21.449	
		8	1	-30.865	9.057	8.085	27.674	121.582	105.769	1347.861
			3	-30.865	8.280	4.863	15.381	85.855	36.796	404.870
	5		-30.865	8.044	3.526	11.869	75.668	19.522	224.123	
	8	10	-30.865	8.125	2.536	7.890	71.494	9.967	98.496	
		2	1	-26.780	12.067	12.067	12.275	248.172	248.172	272.296
			3	-26.780	7.698	7.698	6.889	93.040	93.040	75.998
	5		-26.780	6.601	6.601	5.072	66.846	66.846	39.553	
	2	10	-26.780	5.781	5.781	3.857	47.955	47.955	22.876	
8		1	-30.865	34.556	30.313	28.662	1266.576	1058.689	1518.343	
		3	-30.865	33.742	28.917	14.216	1161.546	882.219	341.538	
	5	-30.865	33.993	28.983	11.307	1169.549	866.469	204.729		
8	10	-30.865	33.858	28.888	8.140	1153.220	847.927	106.282		
	2	1	-26.780	15.622	15.622	11.861	390.349	390.349	231.468	
		3	-26.780	12.878	12.878	6.984	231.040	231.040	78.199	
5		-26.780	11.946	11.946	5.106	184.497	184.497	42.616		
2	10	-26.780	12.066	12.066	3.646	167.718	167.718	20.788		
	8	1	-30.865	67.194	67.826	27.440	4603.887	4782.310	1398.162	
		3	-30.865	66.939	67.699	15.029	4508.284	4638.895	387.225	
5		-30.865	66.675	67.477	11.203	4462.200	4587.222	199.809		
8	10	-30.865	66.688	67.328	8.195	4456.398	4551.466	106.420		
	2	1	-26.780	22.597	22.597	12.162	762.129	762.129	244.327	
		3	-26.780	20.320	20.320	6.716	503.553	503.553	72.748	
5		-26.780	19.418	19.418	5.204	430.610	430.610	43.640		
2	10	-26.780	20.115	20.115	3.599	430.218	430.218	20.391		
	8	1	-30.865	108.579	118.317	26.869	11902.528	14259.553	1344.294	
		3	-30.865	108.185	117.445	15.289	11742.232	13877.573	375.027	
5		-30.865	108.328	117.403	11.473	11757.889	13835.222	214.460		
8	10	-30.865	108.571	118.043	7.979	11799.580	13960.016	102.759		
	2	1	-26.780	31.420	31.420	11.927	1297.693	1297.693	249.215	
		3	-26.780	30.005	30.005	6.688	1008.487	1008.487	71.590	
5		-26.780	30.104	30.104	5.017	972.364	972.364	39.530		
2	10	-26.780	29.547	29.547	3.566	906.943	906.943	19.970		
	8	1	-30.865	164.055	187.416	27.132	27086.351	35529.966	1257.582	
		3	-30.865	163.392	186.002	15.000	26748.599	34712.731	364.521	
5		-30.865	163.930	186.676	11.684	26902.365	34921.680	215.341		
8	10	-30.865	163.562	186.035	8.031	26768.157	34646.567	106.997		

Note. Results over 1000 repetitions. Subscript refers to: (1) TWFE estimator, (2) Wooldridge [2022] estimator, and (3) our estimator. The lowest value across estimators is in bold.

Figure 4b and Table 4 present results from the Poisson DGP, where our estimator performs even better relative to the TWFE and the Wooldridge [2022] ones.

9 Conclusion

We establish identifying assumptions and estimation procedures for the ATT without interference in a DiD setting with staggered treatment adoption and spillovers. Aside from the canonical DiD assumptions, identification requires that once a unit receives treatment, it is no longer influenced by the spillover effect. This means the unit forfeits any spillovers it may have previously received and remains unaffected by spillovers from subsequently treated groups. This assumption, which is likely to hold in many contexts, unifies the multiple definitions of the ATT, simplifying policy evaluation and aligning with the definition of ATT under SUTVA.

To estimate the ATT, we extend the TWFE model approach of Wooldridge [2022] to account for spillovers in linear and non-linear settings. In the case of a balanced panel, our approach can be used to easily calculate the ATT’s standard error. We then revisit Gonzalez-Navarro [2013], who studied the effects of installing an auto theft prevention device known as Lojack. Our correction leads to a slightly larger effect of the policy relative to the original contribution’s specification.

Finally, our Monte Carlo analysis brings attention to the inherent bias-variance trade-off involved in addressing staggered treatment and especially spillovers. We compare three different estimators: the traditional TWFE estimator, which overlooks both staggered adoption and spillovers; the estimator of Wooldridge [2022], which considers staggered adoption but not spillovers; and our proposed estimator, which addresses both factors. Our estimator proves to be competitive in various scenarios.

References

- Tobias Berg, Markus Reisinger, and Daniel Streitz. Spillover effects in empirical corporate finance. *Journal of Financial Economics*, 2021.
- Kirill Borusyak, Xavier Jaravel, and Jann Spiess. Revisiting event study designs: Robust and efficient estimation. Technical report, arXiv.org, 2021.
- Kyle Butts. Difference-in-Differences Estimation with Spatial Spillovers. Papers 2105.03737, arXiv.org, June 2023. URL <https://ideas.repec.org/p/arx/papers/2105.03737.html>.

- Brantly Callaway and Pedro H. C. Sant’Anna. Difference-in-differences with multiple time periods. *Journal of Econometrics*, 2020.
- Damian Clarke. Estimating Difference-in-Differences in the Presence of Spillovers. MPRA Paper 81604, University Library of Munich, Germany, September 2017.
- Clément de Chaisemartin and Xavier D’Haultfoeuille. Two-way fixed effects estimators with heterogeneous treatment effects. *American Economic Review*, 110(9):2964–96, 2020.
- Clément de Chaisemartin and Xavier D’Haultfoeuille. Two-way fixed effects and differences-in-differences with heterogeneous treatment effects: a survey. *The Econometrics Journal*, 2021.
- Marco Gonzalez-Navarro. Deterrence and geographical externalities in auto theft. *American Economic Journal: Applied Economics*, 5(4):92–110, 2013.
- Andrew Goodman-Bacon. Difference-in-differences with variation in treatment timing. *Journal of Econometrics*, 2021.
- Martin Huber and Andreas Steinmayr. A Framework for Separating Individual-Level Treatment Effects From Spillover Effects. *Journal of Business & Economic Statistics*, 39(2):422–436, March 2021.
- László Mátyás and Patrick Sevestre. *The econometrics of panel data: fundamentals and recent developments in theory and practice*. Springer, 2008.
- Robert Minton and Casey B Mulligan. Difference-in-differences in the marketplace. Technical report, National Bureau of Economic Research, 2024.
- Jonathan Roth, Pedro HC Sant’Anna, Alyssa Bilinski, and John Poe. What’s trending in difference-in-differences? a synthesis of the recent econometrics literature. *Journal of Econometrics*, 2023.
- Liyang Sun and Sarah Abraham. Estimating dynamic treatment effects in event studies with heterogeneous treatment effects. *Journal of Econometrics*, 2020.
- Jeffrey M Wooldridge. Two-way fixed effects, the two-way mundlak regression, and difference-in-differences estimators, 2022.
- Jeffrey M Wooldridge. Simple approaches to nonlinear difference-in-differences with panel data. *The Econometrics Journal*, 26(3):C31–C66, 2023.

A Proofs

A.1 Proof of Theorem 1

Under Assumptions 2 and 3, for each group g at time t , we can express Y_{igt} as

$$\begin{aligned} Y_{igt} &= Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) \\ &= Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t) + [Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)] + [Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)] \\ &= \alpha_{ig} + \delta_t + [Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)] + [Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)] + \varepsilon_{igt}, \end{aligned}$$

where the last equality follows from the alternative representation of Assumption 3 given in Equation (4), in which $\mathbb{E}(\varepsilon_{igt}) = 0$ for every group g at time t . Define

$$\begin{aligned} \beta_{igt} &= Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t), \\ \gamma_{igt} &= Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t). \end{aligned}$$

We can then simplify the expression for Y_{igt} as

$$Y_{igt} = \alpha_{ig} + \delta_t + \beta_{igt} + \gamma_{igt} + \varepsilon_{igt}.$$

In this expression, the parameter of interest $ATT_0(g, t)$ for $t \geq g$ is given by

$$ATT_0(g, t) = \mathbb{E}(\beta_{igt} | G_i = g),$$

and $AST(g, t)$ for $t \geq g$ is given by

$$AST(g, t) = \mathbb{E}(\gamma_{igt} | G_i = g).$$

Using these expressions, for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time and $t \geq g$, we can write the expectation of Y_{igt} as

$$\mathbb{E}(Y_{igt}) = \mathbb{E}(\alpha_{ig}) + \delta_t + ATT_0(g, t) + AST(g, t), \quad (9)$$

where we used $\mathbb{E}(\varepsilon_{it}) = 0$.

Now we show that $ATT_0(g, t)$ is identified if and only if $\delta_t + AST(g, t)$ is identified, for every group $g \in \mathcal{G}$ such that $2 \leq g < \infty$ and time $t \geq g$. First, suppose that $\delta_t + AST(g, t)$ is identified. Let d_0 be the identified value. Then we can rewrite Equation (9) as

$$\mathbb{E}(Y_{igt}) = \mathbb{E}(\alpha_{ig}) + d_0 + ATT_0(g, t). \quad (10)$$

Now we show that $\mathbb{E}(\alpha_{ig})$ is identified from the data at $t = 1$. Note first that, under the assumptions of Theorem 1, all units are untreated at $t = 1$. This implies that

$$Y_{ig1} = Y_{ig1}(d_g^1, \mathbf{d}_{(i,g)}^1) = Y_{ig1}(0^1, \mathbf{0}_{(i,g)}^1) = \alpha_{ig} + \delta_1 + \varepsilon_{it},$$

where the last equality follows from Equation (4). Then it follows that

$$\mathbb{E}(Y_{ig1}) = \mathbb{E}(\alpha_{ig} + \delta_1 + \varepsilon_{it}) = \mathbb{E}(\alpha_{ig}), \quad (11)$$

where $\delta_1 = 0$ by Assumption 3 and $\mathbb{E}(\varepsilon_{it}) = 0$ as defined in Equation (4). We can then rewrite Equation (10) as

$$ATT_0(g, t) = \mathbb{E}(Y_{igt}) - d_0 - \mathbb{E}(Y_{ig1}),$$

which shows that $ATT_0(g, t)$ is identified because $\mathbb{E}(Y_{igt})$ and $\mathbb{E}(Y_{ig1})$ are identifiable whenever $g \in \mathcal{G}$, i.e., whenever the group is present in the data.

Conversely, suppose that $ATT_0(g, t)$ is identified. Let b_0 be the identified value. Then we can rewrite Equation (9) as

$$\mathbb{E}(Y_{igt}) = \mathbb{E}(\alpha_{ig}) + \delta_t + b_0 + AST(g, t).$$

Using Equation (11), we can write

$$\delta_t + AST(g, t) = \mathbb{E}(Y_{igt}) - b_0 - \mathbb{E}(Y_{ig1}),$$

which shows that $\delta_t + AST(g, t)$ is identified. ■

A.2 Proof of Theorem 2

By Theorem 1, it suffices to show that $\delta_t + AST(g, t)$ is identified for every $t \geq 2$ under the assumptions of Theorem 2. Note first that Assumption 4 implies $AST(g, t) = 0$. In addition, Assumption 5 states that Λ_∞^0 has a positive measure, implying that the following quantity is identifiable for every $t \geq 2$:

$$\begin{aligned} & \mathbb{E}(Y_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) - \mathbb{E}(Y_{i\infty 1} | (i, \infty) \in \Lambda_\infty^0) \\ &= \mathbb{E}(Y_{i\infty t}(0^t, \mathbf{d}_{(i,g)}^t) | (i, \infty) \in \Lambda_\infty^0) - \mathbb{E}(Y_{i\infty 1}(0^1, \mathbf{0}_{(i,g)}^1) | (i, \infty) \in \Lambda_\infty^0) \\ &= \mathbb{E}(\alpha_{i\infty} | (i, \infty) \in \Lambda_\infty^0) + \delta_t - \mathbb{E}(\alpha_{i\infty} | (i, \infty) \in \Lambda_\infty^0) = \delta_t, \end{aligned}$$

where the equalities follow by Equation (5). This implies that δ_t is identified, which implies that $\delta_t + AST(g, t)$ is identified because $\delta_t + AST(g, t) = \delta_t + 0 = \delta_t$. ■

A.3 Proof of Theorem 3

As in the proof of Theorem 1, under Assumptions 2 and 3, for each group g at time t , we can express Y_{igt} as

$$\begin{aligned} Y_{igt} &= Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) \\ &= Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t) + [Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)] + [Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)] \\ &= \alpha_{ig} + \delta_t + [Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)] + [Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)] + \varepsilon_{igt}. \end{aligned}$$

where $\mathbb{E}(\varepsilon_{igt}) = 0$ for every group g at time t . Define

$$\begin{aligned} \beta_{igt} &= Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t), \\ \gamma_{igt} &= Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t). \end{aligned}$$

We can then simplify the expression for Y_{igt} as

$$Y_{igt} = \alpha_{ig} + \delta_t + \beta_{igt} + \gamma_{igt} + \varepsilon_{igt}.$$

For the group ∞ , this expression simplifies to

$$Y_{i\infty t} = \alpha_{i\infty} + \delta_t + \gamma_{i\infty t} + \varepsilon_{i\infty t}, \quad (12)$$

because $\beta_{i\infty t} = 0$. In addition, Equation (5) states that

$$\begin{aligned} \mathbb{E}(Y_{i\infty t}(0^t, \mathbf{d}_{(i,\infty)}^t) | (i, \infty) \in \Lambda_\infty^0) &= \mathbb{E}(Y_{i\infty t}(0^t, \mathbf{0}_{(i,\infty)}^t) | (i, \infty) \in \Lambda_\infty^0) \\ &= \mathbb{E}(\alpha_{i\infty} | (i, \infty) \in \Lambda_\infty^0) + \delta_t. \end{aligned} \quad (13)$$

The first equality of Equation (13) and the definition of γ_{igt} implies that

$$\mathbb{E}(\gamma_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) = \mathbb{E}(Y_{i\infty t}(0^t, \mathbf{d}_{(i,\infty)}^t) - Y_{i\infty t}(0^t, \mathbf{0}_{(i,\infty)}^t) | (i, \infty) \in \Lambda_\infty^0) = 0.$$

Building on this finding, the second equality of Equation (13), combined with Equation (12), implies that

$$\begin{aligned} \mathbb{E}(\varepsilon_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) &= \mathbb{E}(Y_{i\infty t} - \alpha_{i\infty} - \delta_t | (i, \infty) \in \Lambda_\infty^0) \\ &= \mathbb{E}(Y_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) - \mathbb{E}(\alpha_{i\infty} | (i, \infty) \in \Lambda_\infty^0) - \delta_t \\ &= \mathbb{E}(Y_{i\infty t}(0^t, \mathbf{d}_{(i,\infty)}^t) | (i, \infty) \in \Lambda_\infty^0) - \mathbb{E}(\alpha_{i\infty} | (i, \infty) \in \Lambda_\infty^0) - \delta_t = 0. \end{aligned}$$

Therefore, it follows that $\mathbb{E}(\gamma_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) = \mathbb{E}(\varepsilon_{i\infty t} | (i, \infty) \in \Lambda_\infty^0) = 0$. In addition, since $\mathbb{E}(\varepsilon_{i\infty t}) = 0$, it follows that

$$\mathbb{E}(\varepsilon_{i\infty t} | (i, \infty) \in \Lambda_\infty - \Lambda_\infty^0) = 0,$$

which then implies that $\mathbb{E}(\varepsilon_{igt}) = 0$ for every \mathbf{g} in the extended set of groups $\tilde{\mathcal{G}} \subseteq \{q, \dots, T, (\infty, 0), (\infty, 1)\}$ at time t . Consequently, we can express Y_{igt} for any extended group \mathbf{g} at time t as

$$Y_{igt} = \alpha_{ig} + \delta_t + \beta_{igt} + \gamma_{igt} + \varepsilon_{igt},$$

where $\mathbb{E}(\varepsilon_{igt}) = 0$ for every $\mathbf{g} \in \tilde{\mathcal{G}}$ and $\mathbb{E}(\gamma_{igt}) = 0$ if $\mathbf{g} = (\infty, 0)$.

Now, for every $\mathbf{g} \in \{q, \dots, T\}$, we can write

$$\begin{aligned} Y_{igt} &= \alpha_{ig} + \delta_t + \varepsilon_{ig1} && \text{for } 1 \leq t < q, \\ Y_{igt} &= \alpha_{ig} + \delta_t + \gamma_{igt} + \varepsilon_{igt} && \text{for } q \leq t < \mathbf{g}, \\ Y_{igt} &= \alpha_{ig} + \delta_t + \beta_{igt} + \varepsilon_{igt} && \text{for } \mathbf{g} \leq t \leq T, \end{aligned}$$

These expressions are obtained by the following arguments:

- The first expression is obtained by recognizing that, in the pre-treatment periods, neither the treatment effect nor the spillover effects are present, represented by $\beta_{igt} = 0$ and $\gamma_{igt} = 0$.
- The second expression is obtained by recognizing that, in the post-treatment periods where units in group \mathbf{g} are not yet treated, there is no treatment effect ($\beta_{igt} = 0$), while spillover effects may occur, represented by γ_{igt} .
- The third expression is obtained by recognizing that, in the post-treatment periods where units in group \mathbf{g} have been treated, the treatment effect may present, represented by β_{igt} , but units are not subject to spillover effects by Assumption 4, represented by $\gamma_{igt} = 0$.

We can combine these three expressions into one unified expression, encompassing every group $\mathbf{g} \in \{q, \dots, T\}$ at every period $1 \leq t \leq T$, as follows:

$$Y_{igt} = \alpha_{ig} + \delta_t + \sum_{t'=\mathbf{g}}^T \beta_{igt'} \mathbf{1}(t = t') + \sum_{t'=q}^{\mathbf{g}-1} \gamma_{igt'} \mathbf{1}(t = t') + \varepsilon_{igt},$$

which we can further write as

$$Y_{igt} = \alpha_{ig} + \delta_t + \sum_{t'=\mathbf{g}}^T \beta_{igt'} \mathbf{1}(t = t') D_{igt} + \sum_{t'=q}^{\mathbf{g}-1} \gamma_{igt'} \mathbf{1}(t = t') S_{igt} + \varepsilon_{igt}, \quad (14)$$

since $D_{igt} = 1$ for $t \geq \mathbf{g}$ and $S_{igt} = 1$ for $q \leq t < \mathbf{g}$ for each $\mathbf{g} \in \{q, \dots, T\}$ according to their definitions.

Next, for groups $\mathbf{g} \in \{(\infty, 0), (\infty, 1)\}$, we can write

$$\begin{aligned} Y_{i,(\infty,0),t} &= \alpha_{i,(\infty,0)} + \delta_t + \varepsilon_{i,(\infty,0),t} && \text{for } 1 \leq t \leq T, \\ Y_{i,(\infty,1),t} &= \alpha_{i,(\infty,1)} + \delta_t + \varepsilon_{i,(\infty,1),t} && \text{for } 1 \leq t < q, \\ Y_{i,(\infty,1),t} &= \alpha_{i,(\infty,1)} + \delta_t + \gamma_{i,(\infty,1),t} + \varepsilon_{i,(\infty,1),t} && \text{for } q \leq t \leq T, \end{aligned}$$

These can also be unified similarly to Equation (14), where we write Y_{igt} for every $\mathbf{g} \in \{(\infty, 0), (\infty, 1)\}$ at every period $1 \leq t \leq T$ as

$$Y_{igt} = \alpha_{ig} + \delta_t + \sum_{t'=\mathbf{g}}^T \beta_{igt'} \mathbf{1}(t=t') D_{igt} + \sum_{t'=q}^{\mathbf{g}-1} \gamma_{igt'} \mathbf{1}(t=t') S_{igt} + \varepsilon_{igt}, \quad (15)$$

where $\sum_{t'=\mathbf{g}}^T$ is considered a null summation, interpreted as $\sum_{t'=\infty}^T$, and $\sum_{t'=q}^{\mathbf{g}-1}$ is interpreted as $\sum_{t'=q}^T$. Note that $D_{igt} = 0$ if $\mathbf{g} \in \{(\infty, 0), (\infty, 1)\}$, reflecting the absence of treatment effects in these groups, and that $S_{igt} = 0$ if $\mathbf{g} \in \Lambda_\infty^0$, indicating no spillover effects for this group. One might be concerned that the β_{igt} and γ_{igt} values corresponding to these cases are not determined, but their definitions imply that $\beta_{igt} = 0$ if $\mathbf{g} \in \{(\infty, 0), (\infty, 1)\}$ and that $\gamma_{igt} = 0$ if $\mathbf{g} \in \Lambda_\infty^0$.

Now, we can combine Equation (14) and Equation (15) and write Y_{igt} for any extended group $\mathbf{g} \in \tilde{\mathcal{G}}$ at any time period $1 \leq t \leq T$ as

$$\begin{aligned} Y_{igt} &= \sum_{\mathbf{g}' \in \tilde{\mathcal{G}}} \mathbf{1}(\mathbf{g} = \mathbf{g}') \left(\alpha_{ig} + \delta_t + \sum_{t'=\mathbf{g}'}^T \beta_{ig't'} \mathbf{1}(t=t') D_{igt} + \sum_{t'=q}^{\mathbf{g}'-1} \gamma_{ig't'} \mathbf{1}(t=t') S_{igt} + \varepsilon_{igt} \right) \\ &= \alpha_{ig} + \delta_t + \sum_{\mathbf{g}' \in \tilde{\mathcal{G}}} \sum_{t'=\mathbf{g}}^T \beta_{ig't'} \mathbf{1}(\mathbf{g} = \mathbf{g}') \mathbf{1}(t=t') D_{igt} + \sum_{\mathbf{g}' \in \tilde{\mathcal{G}}} \sum_{t'=q}^{\mathbf{g}'-1} \gamma_{ig't'} \mathbf{1}(\mathbf{g} = \mathbf{g}') \mathbf{1}(t=t') S_{igt} + \varepsilon_{igt}. \end{aligned} \quad (16)$$

Next, note that Equation (6) is a pooled regression of the variables that encompass all groups $\mathbf{g} \in \tilde{\mathcal{G}}$. Specifically, let $j = 1, \dots, N$ be the index for units in this pooled regression, where $N \equiv \sum_{\mathbf{g} \in \tilde{\mathcal{G}}} N_{\mathbf{g}}$ is the total number of observations across all groups. Let Y_{jt} be the outcome variable for index j , with D_{jt} and S_{jt} defined similarly. Let \mathbf{X}_{jt} be the vector of regressors in Equation (6), namely indicators of $\mathbf{g} \in \tilde{\mathcal{G}}$, indicators of t , interactions between D_{jt} and indicators of (\mathbf{g}, t) , and interactions between S_{jt} and indicators of (\mathbf{g}, t) .

The estimation of this pooled regression involves computing averages of these variables across $j = 1, \dots, N$. This is equivalent to the weighted averages of (i, \mathbf{g}) , pooled across all groups, where the weights are determined by the relative sizes of the groups in the

dataset. For example, the average of Y_{jt} across $j = 1, \dots, N$ is calculated as

$$\frac{1}{N} \sum_{j=1}^N Y_{jt} = \frac{1}{N} \sum_{\mathbf{g} \in \tilde{\mathcal{G}}} \sum_{i=1}^{N_{\mathbf{g}}} Y_{igt} \equiv \sum_{\mathbf{g} \in \tilde{\mathcal{G}}} w_{\mathbf{g}} \cdot \frac{1}{N_{\mathbf{g}}} \sum_{i=1}^{N_{\mathbf{g}}} Y_{igt},$$

where the weights $w_{\mathbf{g}}$ are defined as $w_{\mathbf{g}} = N_{\mathbf{g}}/N$.

We derive the population regression of Equation (6) as follows. We define the expectation across j as a weighted average of expectations across all groups. Specifically, the expectation of Y_{jt} is defined as

$$\bar{\mathbb{E}}(Y_{jt}) \equiv \sum_{\mathbf{g} \in \tilde{\mathcal{G}}} \bar{w}_{\mathbf{g}} \mathbb{E}(Y_{igt}),$$

where $\bar{w}_{\mathbf{g}}$ is the weight for group \mathbf{g} such that $\sum_{\mathbf{g} \in \tilde{\mathcal{G}}} \bar{w}_{\mathbf{g}} = 1$. We assume that the relative sizes of the groups are fixed both across the sampling processes and in the population, meaning that $\bar{w}_{\mathbf{g}} = w_{\mathbf{g}} = N_{\mathbf{g}}/N$. This corresponds to the asymptotics where $N \rightarrow \infty$ along the sequence $(N, 2N, 3N, 4N, \dots)$ that maintains the relative sizes of the groups. This setting aligns with that of [Borusyak et al. \[2021\]](#).

Now we proceed to prove the theorem. Note first that the interaction terms in $(\mathbf{X}_{j1}, \dots, \mathbf{X}_{jT})$ identify the extended group label \mathbf{g} and vice versa, because the interactions of D_{igt} identifies the group label $g \in \{1, \dots, T\}$ and the interactions of S_{igt} distinguishes $(\infty, 0)$ and $(\infty, 1)$. This implies that

$$\bar{\mathbb{E}}(Y_{jt} | \mathbf{X}_{j1}, \dots, \mathbf{X}_{jT}) = \mathbb{E}(Y_{igt} | \mathbf{X}_{j1}, \dots, \mathbf{X}_{jT}).$$

Then, by Equation (16):

$$\begin{aligned} \mathbb{E}(Y_{jt} | \mathbf{X}_{j1}, \dots, \mathbf{X}_{jT}) &= \mathbb{E}(\alpha_{i\mathbf{g}}) + \delta_t + \sum_{\mathbf{g} \in \tilde{\mathcal{G}}} \sum_{t'=\mathbf{g}}^T \mathbb{E}(\beta_{i\mathbf{g}'t'}) \mathbf{1}(\mathbf{g} = \mathbf{g}') \mathbf{1}(t = t') D_{igt} \\ &\quad + \sum_{\mathbf{g}' \in \tilde{\mathcal{G}}} \sum_{t'=q}^{g-1} \mathbb{E}(\gamma_{i\mathbf{g}'t'}) \mathbf{1}(\mathbf{g} = \mathbf{g}') \mathbf{1}(t = t') S_{igt}. \end{aligned}$$

This shows that the coefficient associated with $\mathbf{1}(\mathbf{g} = \mathbf{g}') \mathbf{1}(t = t') D_{igt}$ for $g \in \{2, \dots, T\}$ at time t is $\mathbb{E}(\beta_{igt})$. Then, by the definition of β_{igt} :

$$\mathbb{E}(\beta_{igt}) = \mathbb{E}(Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)),$$

where the right-hand side is the definition of $ATT_0(g, t)$. ■

A.4 Proof of Corollary 1

Similarly to the proof of Theorem 1, for each group g at time t , we can express Y_{igt} as

$$\begin{aligned} Y_{igt} &= Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) \\ &= Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t) + [Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)] + [Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)]. \end{aligned}$$

Then, under Assumption 3', we can write the expectation of Y_{igt} as

$$\mathbb{E}(Y_{igt}|G_i = g) = \mathbb{E}(\exp\{\alpha_{ig}\}|G_i = g) \cdot \exp\{\delta_t\} + ATT_0(g, t) + AST(g, t),$$

where

$$ATT_0(g, t) = \mathbb{E}(Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t) - Y_{igt}(0^t, \mathbf{0}_{(i,g)}^t)|G_i = g),$$

and

$$AST(g, t) = \mathbb{E}(Y_{igt}(d_g^t, \mathbf{d}_{(i,g)}^t) - Y_{igt}(d_g^t, \mathbf{0}_{(i,g)}^t)|G_i = g).$$

Then, by replicating the arguments in Theorem 1 that starts from Equation (9), it is straightforward to show that $ATT_0(g, t)$ is identified if and only if $\mathbb{E}(\exp\{\alpha_{ig}\}) \cdot \exp\{\delta_t\} + AST(g, t)$ is identified. ■

A.5 Proof of Corollary 2

By Corollary 1, it suffices to show that $\mathbb{E}(\exp\{\alpha_{ig}\}) \cdot \exp\{\delta_t\} + AST(g, t)$ is identified. We proceed by separately identifying the three objects $\mathbb{E}(\exp\{\alpha_{ig}\})$, $\exp\{\delta_t\}$, and $AST(g, t)$. First, Assumption 4 implies that $AST(g, t) = 0$, identifying $AST(g, t)$. Second, consider the units (i, g) that belong to Λ_∞^0 , which are never-treated units that are not affected by spillover effects. For these units, the following moment equality holds for any t under Assumption 3' and assumption 5 [Mátyás and Sevestre, 2008, Chapter 18.3.1]:

$$\mathbb{E} \left(Y_{i\infty t} - \exp\{\delta_t\} \frac{(1/T) \sum_{t'=1}^T Y_{i\infty t'}}{(1/T) \sum_{t'=1}^T \exp\{\delta_{t'}\}} \middle| (i, g) \in \Lambda_\infty^0 \right) = 0 \quad \text{for all } t = 1, \dots, T, \quad (17)$$

which can be verified straightforwardly. Evaluating this moment equality at $t = 1$ yields

$$\mathbb{E} \left(Y_{i\infty 1} - \frac{(1/T) \sum_{t'=1}^T Y_{i\infty t'}}{(1/T) \sum_{t'=1}^T \exp\{\delta_{t'}\}} \middle| (i, g) \in \Lambda_\infty^0 \right) = 0,$$

since $\delta_1 = 0$. This equation identifies the term $(1/T) \sum_{t'=1}^T \exp\{\delta_{t'}\}$. Then, evaluating Equation (17) for $t \geq 2$ identifies $\exp\{\delta_t\}$ for each $t \geq 2$.

Lastly, because all units are untreated at $t = 1$ by the assumption, it follows that

$$\mathbb{E}(Y_{ig1}) = \mathbb{E}(Y_{ig1}(0, \mathbf{0}_{(i,g)}^1)) = \mathbb{E}(\exp\{\alpha_{ig}\}).$$

This implies that $\mathbb{E}(\exp\{\alpha_{ig}\})$ is identified for every $g \in \mathcal{G}$, because Y_{ig1} is identifiable whenever $g \in \mathcal{G}$, i.e., whenever the group is present in the data. ■